1. F.S-2009

Nº 007471

B-JGT-J-DIA

CHEMISTRY

Paper I

Time Allowed: Three Hours

Maximum Marks : 200

INSTRUCTIONS

Candidates should attempt question I and 5 which are compulsory, and any THREE of the remaining questions, selecting at least ONE question from each Section.

The number of marks carried by each question is indicated at the end of the question.

Assume suitable data if considered necessary, and indicate the same clearly.

Answers must be written in ENGLISH.

Unless otherwise indicated, symbols and notations have their usual meanings.

(Planck's constant $h = 6.62 \times 10^{-34} \text{ Js}$

Boltzmann constant $k = 1.38 \times 10^{-23} \text{ J K}^{-1}$

Electron rest mass $m = 9.1 \times 10^{-28} g$

SECTION A

- 1. Answer any four of the following:
 - (a) Show that free energy change $-\Delta G_{P,T}$ is the total amount of non-mechanical work.
 - (b) Give a short account of "liquid crystals". 10
 - (c) Given the following $E_{Ag, AgBr}^{\circ} = -0.073 \text{ V}$, K for the reaction $Ag + \frac{1}{2} Br_2 \rightleftharpoons AgBr$ is 6.46×10^{16} at 25° C, calculate $E_{Br_2, Br}^{\circ}$. Write the suitable chemical cell.
 - (d) Explain the Graham's of effusion. 10
 - (e) Calculate the freezing temperature of water when the pressure's increased by two atm. $(\Delta H_f = 79.9 \text{ cal A}^{-1}; \ \rho_{ice} = 0.92 \text{ g cc}^{-1}; \\ \rho_{Hg} = 13.6 \text{ g cc}^{-1}.)$
- 2. (a) Explain the Maxwell's distribution of molecular velocities and show that

$$dn_c / dc = 4 \pi n \left(\frac{m}{2\pi kT}\right)^{3/2} c^2 e^{-\frac{mc^2}{2kT}}$$

where 'n' is the number of molecules (of mass 'm')
per cc, dn_c/dc is the number of molecules per
unit range of velocity, 'c' is the velocity and the
rest of the symbols have their usual significance. 20

10

(b) Define a unimolecular reaction. Explain unimolecular reaction with the help of collision theory. Guess the order of the following reaction $A + B \rightarrow 2C + D$.

20

3. Discuss the "Uncertainty principle" in detail. Show that for a macroscopic body this principle is of no importance. What was Einstein's opinion about this principle? Discuss.

40

4. (a) Show that $H_2 + Br_2 \rightarrow 2$ HBr does not have an order of reaction and the formation of HBr inhibits the reaction.

30

(b) A first order decomposition reaction is characterised by a rate constant $4 \times 10^{-2} \text{ min}^{-1}$. Find out the time required to complete one-third of the reaction.

10

SECTION B

5.	Answer any four of the following:		
	(a)	Draw electronic configuration, atomic and molecular orbitals of F_2 .	10
-	(b)	Draw the structure of (i) dichromate ion (ii) solid HF.	10
	(c)	Explain the 'partition function'. Deduce the 'internal energy' and 'entropy' in terms of partition function.	10
	(d)	Explain how the filling of 4f energy level affects the rest of the periodic table.	10
	(e)	What is 'relaxation time'? Explain the 'relaxation method' in the study of late of reaction.	10
6.	(a)	Draw the shape of various 'd' orbitals and explain as to why they are split in two groups e_g and t_{2g} in an outshedral ligand field.	20
	(b)	How do cryoscopic and molar conductivity measurements help in establishing the structure of complexes? Explain with examples.	20
7.	(a)	Discuss the laws of photochemistry. What is meant by "One Einstein"?	
		On irradiation with light of 300 nm , propionaldehyde yielded CO. If the rate of formation of CO is $2.04 \times 10^{-9} \text{ moles s}^{-1}$ and intensity of light is $10,000 \text{ ergs s}^{-1}$, calculate the quantum yield of CO.	30

(b) 5% of energy of a 200 W incandescent lamp emits visible light ($\lambda = 600$ nm). How many quanta of light is emitted per second?

10

20

- 8. (a) What is Born Haber cycle? Using a modified form of this cycle explain why Li is as strong a reducing agent as Cs.
 - (b) Discuss limitations and practical uses of liquid ammonia as non-aqueous solvent. What happens when ¹⁵NH₄Cl is dissolved in unfacelled liquid NH₃ and the solvent evaporated? 20

www.sudy cidelindia.com