
1. Following two wave trains are approaching each other. $y_1 = a \sin 200 \pi t$ $y_2 = a \sin 208 \pi t$ The number of beats heard per second is :				
A. 8	B. 4	C. 1	D. 0	
2. One of the geo-static	onary satellites of India is	s vertically above		
A. New Delhi	B. Mumbai	C. Allahabad	D. None of these	
3. Light of wavelength equal to	$2400 \ge 10^{-10} $ m in air wil	ll become light of wavel	ength in glass ($\mu = 1.5$)	
-	B. 7200 x 10 ⁻¹⁰ m	C. 1080 x 10 ⁻¹⁰ m	D. none of these	
	ry to primary turns is 4:5 all losses) to power input		at will be the ratio of	
A. 4:9	B. 9:4	C. 5:4	D. 1:1	
5. Lenz's law applies to A. electrostatics)	B. lenses		
C. electro-magnetic inc	luction	D. cinema slides		
6. If a proton and anti-preleased ?	proton come close to each	h other and annihilate, h	ow much energy will be	
A. 1.5 x 10 ⁻¹⁰ J	B. 3 x 10 ⁻¹⁰ J	C. 4.5 x 10 ⁻¹⁰ J	D. none of these	
-	s, what will be the result	t		
?	C intrinsis			
A. <i>n</i> -type B. <i>p</i> -type semi- semi-	C. intrinsic D. none of semi-			
conductor conductor	conductor these			
8. A charge is placed at the centre of a cube, what is the electric flux passing through one of its faces?				
A. $(1/6) \ge (q/\epsilon_0)$	B. q/ε_0	C. $6q/\varepsilon_0$	D. None of these	
9. What is the degree of freedom in case of a mono atomic gas ?				
A. 1	B. 3	C. 5	D. None of these	
10. The ratio of secondary to primary turns is 4:5. If power input is <i>P</i> , what will be the ratio of power output (neglect all losses) to power input ?				



11. Speed of recession of galaxy is proportional to its distance

A. directly	B. inversely	C. exponentially	D. none of these		
12. If a substance goes in a magnetic field and is pushed out of it, what is it ?A. ParamagneticB. FerromagneticC. DiamagneticD. Antiferromagnetic					
	Ū.				
13. Which is not a scal	ar quantity?				
A. Work	B. Power	C. Torque	D. Gravitational Constant		
14. Minimum energy r A13.6 eV	equired to excite an elect B. 13.6 eV	tron in a Hydrogen atom C. 10.2 eV	in ground state is : D. 3.4 eV		
15. If Gravitational Co satellite orbiting aroun	nstant is decreasing in ti d earth ?	me, what will remain un	changed in case of a		
A. Time period	B. Orbiting radius	C. Tangential velocity	D. Angular velocity		
16. If a transparent medium of refractive index $\mu = 1.5$ and thickness $t = 2.5 \times 10^{-5}$ m is inserted in front of one of the slits of Young's Double Slit experiment, how much will be the shift in the interference pattern? The distance between the slits is 5.0×10^{-3} cm and that between slits and screen is 100 cm.					
A. 5 cm	B. 2.5 cm	C. 0.25 cm	D. 0.1 cm		
17. How does light pro	pagate in optical fibres?				
A. Total internal reflection	B. Refraction	C. Reflection	D. None of these		
18. Dispersion of light	is due to				
A. wavelength	B. intensity of light wing conclusions is corre	C. density of medium ect regarding	D. none of these		
A. No force is acting o	n the body				
Ũ	s acing on the body is ze	ro			
C. The body is in vacu	-				
D. The forces acting on the body do not constitute a couple					
20. Energy released in	20. Energy released in stars is due to				
A. Fission	B. Fusion	C. Combustion	D. Chemical reaction		
21. 13 days is the half- 1/16th of the original s	life period of a sample. A ubstance ?	After how many days, the	e sample will become		
A. 52	B. 3.8	C. 3	D. none of these		
22 Alter lete	- 4 4 1 * - 1				

22. Absolute zero is the temperature at which

A. water solidifies B. all gases become liquid C. motion of molecules becomes minimum D. everything solidifies 23. Motion of liquid in a tube is described by D. Archimedes' A. Bernaulli's Theorem B. Poiseuille Equation C. Stoke's Law Principle 24. Molecular motion shows itself as A. Temperature B. Internal Energy C. Friction **D.** Viscosity 25. Which is this gate ? A. AND **B. NAND** C. OR D. NOR 26. Energy bands in solids are a consequence of A. Ohm's Law **B.** Pauli's Exclusion Principle D. Heissenberg's Uncertainty Principle C. Bohr's Theory 27. A boy of mass M stands on the floor of an elevator moving downwards with an acceleration a which is less than g. The force exerted by the boy on the floor of the elevator is C. Mg – Ma A. Mg x Ma B. g + aD.Mg + Ma28. A body A of mass m₁ exerts a force on another body B of mass m₂. If the acceleration of B be a₂, then the acceleration (in magnitude) of A is A. m_2/m_1 (a₂) B. $m_1m_2 a_2$ C. m_1/m_2 (a₂) D. $(m_1 + m_2) a_2$ 29. What does not change when sound enters from one medium to another ? A. Wavelength B. Speed C. Frequency D. none of these 30. Resolving power of a microscope depends upon A. wavelength of light used, directly B. wavelength of light used, inversely C. frequency of light used D. focal length of objective 31. An astronaut of weight Mg is in a rocket accelerating upward with an acceleration of 4g. The apparent weight of the astronaut will be A. 5Kg C. Mg B. 4Kg D. zero 32. One proton beam enters a magnetic field of 10^{-4} m/s normally, sp. charge = 10^{11} C/kg, velocity = 10^9 m/s. What is the radius of the circle describe by it ? A. 0.1 m B. 100 m C. 10 m D. none of these 33. If a black body radiates 20 calories per second at 227°C, it will radiate at 727°C

A. 10 calories per second	B. 80 calories per second	C. 320 calories per second	D. none of these		
34. If a carnot engine is working with source temperature equal to 227°C and its sink temperature is at 27°C, its efficiency will be					
A. 20%	B. 10%	C. 67%	D. 50%		
35. If the frequency of a energy is	an oscillating particle is <i>r</i>	<i>i</i> , then the frequency of c	oscillation of its potential		
A. n	B. 2n	C. n/2	D. 4n		
A. X-rays C. Infra-red rays	ates at a frequency of 1 C	GHz, it gives : B. Micro-waves D. None of these			
37. Earth's atmosphere					
A. Ultra-violet rays	B. Infra-red rays	C. X-rays	D. Micro-waves		
38. Cathode rays consis	st of				
A. Photons	B. Electrons	C. Protons	D. α -particles		
	is moving with a velocity edded. At the point of co		e system		
1:2. The smaller part be	ing with velocity V in spacecomes stationary. What	is the velocity of the othe	er part ?		
A. 4V	B. V	C. 4V/3	D. 2V/3		
41. A thief steals a box of weight W & jumps from the third floor of a building. During jump, he experiences a weight of					
A. W	B. 3W	C. 1.5W	D. zero		
42. Two electron beams are moving parallel in space but in opposite directions; thenA. they will attract each otherB. they will repel each otherC. no interaction will take placeD. none of these					
43. Two wires with resi 2R and R is	istances R and 3R are con	nnected in parallel, the ra	tio of heat generated in		
A. 1 : 3	B. 2 : 1	C. 1 : 4	D. 4 : 1		

44. A wire is drawn such that its radius changes from r to 2r, the new resistance is

A. 2 times	B. 4 times	C. 8 times	D. 1/16 times			
45. In solids, inter-ator A. totally repulsive C. combination of (a) a		B. totally attractive D. none of these				
A. he is taken abackB. he is afraidC. due to inertia of rest						
47. What should be the the string just does not	e minimum velocity at the slack ?	e highest point of a body	tied to a string, so that			
A. $\sqrt{(Rg)}$	B. $\sqrt{(5Rg)}$	C. $(R/g)^{3/2}$	D. $\sqrt{(2Rg)}$			
48. If a person standingA. increaseC. remain same	g on a rotating disc stretc	hes out his hands, the sp B. decrease D. none of these	eed will:			
49. EMF is most close A. mechanical force	ly related to B. potential difference	C. electric field	D. magnetic field			
50. Planetary system in A. conservation of ene C. conservation of ang		bes B. conservation of lines D. none of these	ar momentum			
51. Lenz's law is based A. energy	l upon B. momentum	C. angular momentum	D. inertia			
52. Faraday's second la	aw states that mass depos	sited on the electrode is d	lirectly proportional to			
A. atomic mass	B. atomic mass x velocity	C. atomic mass/valency	y D. valency			
53. Unit of power is A. kilowatt hour	B. kilowatt per hour	C. kilowatt	D. erg			
54. Power can be expre A. F.v	essed as B. $1/2$ (Fv ²)	C. F.t	D. F x v			
55. Units of coefficient A. Nms ⁻¹	t of viscosity are B. Nm ² s ⁻¹	C. Nm^{-2} s	D. Nms ⁻²			

56. Dimensions of torq A. MLT ⁻²	ue are B. ML^2T^{-2}	$C. M^2 L^2 T^{-2}$	D. $ML^{-2}T^{-2}$	
57. A body of weight <i>n</i> extending the string is	ng is hanging on a string.	, which extends its lengt	h by <i>l</i> . The work done in	
A. mg l	B. mg l/2	C. 2 mg l	D. none of these	
58. The water droplets	in free fall are spherical	due to		
A. gravity	B. viscosity	C. surface tension	D. inter-molecular attraction	
59. A ball of mass 1Kg	s is accelerating at a rate	of 1ms ⁻² . The rate of cha	inge of momentum is	
A. 1 Kg ms ⁻²	B. 2 Kg ms ⁻²	C. 3 Kg ms ⁻²	D. 4 Kg ms ^{-2}	
60. A body orbitting ar orbit of a satellite. The	ound earth at a mean rad period of the body is	lius which is two times a	s great as the parking	
A. 4 days	B. $2\sqrt{2}$ days	C. 16 days	D. 64 days	
61. Gamma rays are		.0.		
A. high energy electron	18	B. low energy electrons	5	
C. high energy electro-	magnetic waves	D. high energy positror	15	
62. Which is the most a	abundant metal in the ear	rth's crust?		
A. Fe	B. Al	C. Ca	D. Na	
63. Which one does no	t give a precipitate with	excess of NaOH?		
A. ZnSO ₄	B. FeSO ₄	C. AgNO ₃	D. HgCl ₂	
64. What volume of CO_2 will be liberated at NTP of 12 gm of carbon is burnt in excess of oxygen?				
A. 11.2 litres	B. 22.4 litres	C. 2.24 litres	D. 1.12 litres	
65. Which base is found only in nucleotides of RNA?				
A. Adenine	B. Uracil	C. Guanine	D. Cytosine	
66. Ascorbic acid is the	e chemical name of			
A. Vitamin B_6	B. Vitamin A	C. Vitamin C	D. Vitamin D	

67. A hydrocarbon has carbon and hydrogen. Its molecular weight is 28. Its possible formula would be

A. C ₃ H ₆	B. C_2H_4	C. CH ₄	D. C ₄ H ₈	
68. The first Noble Pri A. Faraday	ze in chemistry was give B. Cnrizzaro	n to C. Mendeleevs	D. Moseley	
69. Four different collo action?	oids have the following g	old number. Which one	has its most effective	
A. 10	B. 30	C. 20	D. 40	
70. Which is an examp A. Polythene	le of thermosetting poly B. PVC	mer? C. Neoprene	D. Bakelite	
71. The number of unp A. 3	baired electrons in ferrous B. 2	s ion is C. 4	D. 5	
72. Strongest reducing A. K	agent is B. Mg	C. Al	D. Ba	
73. Which of the follow A. Ra	wing is man-made eleme B. U	nt? C. Np	D. C – 4	
74. Which of the following statements is/are correct? A. Boiling point of alkylhalide is greater than its corresponding alkane B. In water, solubility of $CH_3OH > C_2H_5OH >$ C_6H_5OH C. Aniline is a weaker base than NH_3 D. All of the above				
75. Which amine of the A. Ethylamine	e following will not ansv B. Methylamine	ver Carbylamine reaction C. Dimethylamine	n? D. Phenylamine	
76. Tollen's reagent ca A. (CH ₃) ₂ – CHOH	n be used to detect B. CH ₃ – CO.CH ₃	C. CH ₃ CH ₂ CHO	D. CH ₃ OCH ₃	
77. Glycerol on heating with Potassium bisulphate yieldsA. AcetoneB. GlyceraldehydeC. AcroleinD. Propanol				
78. Salicylic acid on he A. Benzene	eating with sodalime give B. Calcium salicylate		D. Phenol	
79. Which one of the for A. Ethanol	ollowing will not give io B. Ethanal	doform test? C. 2-propanone	D. None of these	

80. The rusting of iron A. Fe	is catalysed by B. O ₂	C. Zn	D. H ⁺		
81. 100 ml of a liquid A was mixed with 25 ml of a liquid B to give non-ideal solution of A-B mixture. The volume of this mixture will be					
A. 75 ml		B. 125 ml exact			
C. fluctuating between 75 ml and 125 ml D. close to 125 ml but not to exceed 125 ml					
	compound having the for				
A. 3, 3 - dimethyl - 1 -		B. 1, 1 - dimethyl - 3 -			
C. 1,1, 1 - dimethyl - 2	2 - propene	D. 3, 3, 3 - dimethyl -	I - I propene		
83. Which of the follow	wing compounds will be	optically active?			
A. (OH ₃) ₂ – CHOH	B. CH ₃ - CH ₂ - CH ₂ - CH ₃	C. CH ₃ – CHCl.COOF	I D. (CH ₃) ₃ .C.Cl		
84. The major compon	ents of brass are				
A. Zn and Sn	B. Cu and Zn	C. Fe and Ni	D. Zn and Fe		
85. Lunar castic is					
A. Silver Chloride	B. Silver Nitrate	C. Sodium Hydroxide	D. Potassium Nitrate		
86. When hot iron is ex	xposed in hot water vapo	our, the compound forme	d is		
A. FeO	B. Fe_2O_4	C. Fe ₃ O ₄	D. Fe_2 (OH) ₂		
87. Which of the follow	wing halide is not oxidise	ed by MnO ₂ ?			
A. F	B. Cl	C. Br	D. I ⁻		
88. The outermost elec	ctronic configuration of t	he most electronegative	element is		
A. $ns^2 np^3$		C. $ns^2 np^5$			
89. Shape of CO_2 is					
A. tetrahedral	B. trigonal	C. bent	D. linear		
90. The catalyst used i A. Al ₂ O ₃	n the manufacture of H ₂ S B. Cr ₂ O ₃	SO ₄ by contact process is C. V ₂ O ₅	D. MnO ₂		
91. The composition o	f the common glass is				
A. Na ₂ O.CaO.6SiO ₂	B. Na ₂ O.Al ₂ O ₃ .2SiO ₂	C. CaO.Al ₂ O ₃ .2SiO ₂	D. Na ₂ O.CaO.Al ₂ O ₃ .6SiO ₂		

92. In a borax lead test A. Chromium	, the brown colour is due B.Cobalt	e to C. Manganese	D. Iron		
93. Which of the follow A. Urea	0	C. Benzene Hexachloride	D. Potassium		
94. Which one of the for Table?	ollowing belongs to repro	esentative group of elem	ents in the Periodic		
A. Lanthanum	B. Argon	C. Chromium	D. Aluminium		
95. Which one of the fe A. Tritium	ollowing is not an isotop B. Deuterium	e of Hydrogen? C. Ortho-hydrogen	D. None of the above		
	96. In the reaction $I_2 + 2S_2O_3^{2^-} = 2I^- + S_4O_6^{2^-}$, equivalent weight of iodine will be equal to A. its molecular weight C. 1/4 the molecular weight D. twice the molecular weight				
97. Which of the following is the most powerful oxidising agent? A. F_2 B. Cl_2 C. Br_2 D. l_2					
98. From the following strongest acid?	g values of dissociating c	onstants of four acids, w	hich value represents the		
A. 2×10^{-2}	B. 0.02 x 10 ⁻¹	C. 3 x 10 ⁻³	D. 2.0 x 10 ⁴		
99. In which of the foll	lowing cases, does the re	action go the farthest for	completion?		
A. $K = 10^3$	B. K = 10^{-2}	C. K = 10	D. K = 1		
100. The reaction whic	ch proceeds in the forwar	d direction is			
A. $Fe_2O_3 + 6HCl \rightarrow 2H$ C. $SnCl_4 + Hg_2Cl_2 \rightarrow S$		B. $NH_3 + H_2O + NaCl$ D. $2CuI + I_2 + 4K^+ \rightarrow$			
101. The substance capable of being drawn into fine wire is called					
A. malleable	B. tensile	C. ductile	D. mild		
102. The idea that mos is given by	t of the mass of an atom	is concentrated in a very	small core, i.e., nucleus		
A. Amedo Avogadro	B. Rutherford	C. Bohr	D. Henery Mosley		
103. Which of the follo A. $N_2H_5^+$	owing does contain a co- B. BaCl ₂	ordinate covalent bond? C. HCl	D. H ₂ O		

104. Which of the following containsA. CCl_4 B. $CaCl_2$	both covalent and ic C. NH4C		D. H ₂ O	
105. Keeping in view the periodic law elements should have the maximum e	1		ich of the following	
A. Oxygen B. Nitrogen	C. Fluori	ne l	D. Astatine	
106. The electronic configuration of e	lement atomic numb	per 37 is		
A. $(2, 8) 3s^2 3p^6 3d^{10} 4s^2 4p^6 5s^1$	B. (2, 8)	3s ² 3p ⁶ 3d ¹⁰ 4s ² 5s	$s^{6}4p^{5}$	
C. $(2, 8) 3s^2 3p^6 4s^2 3d^9 5s^1 4p^5$	D. none of	of these		
107. The pH of 0.1 M solution of a we the acid?	eak acid is 3. What i	s the value of ic	onisation constant for	
A. 0.1 B. 10 ⁻³	C. 10 ⁻⁵	. 0'	D. 10 ⁻⁷	
108. Pure Aniline is a	C human			
A. brown coloured B. colourless	liquid c. blown	coloured	D. colourless solid	
-				
109. Sulphide ores are generally conce	-	na hu sanhan '	D tomasias	
A. roasting B. froth floata	uton C. reduct	ng by carbon	D. tempering	
110. One mole of CO_2 contains				
A. 6.02×10^{23} atoms of C		10^{23} atoms of (С	
C. 18.1 x 10^{23} molecules of CO ₂	D. 3 gm	atom of CO ₂		
C [×]				
111. The Avogadro Number or a mole	-	22		
A. $6.02 \ge 10^{23}$ ions B. $6.02 \ge 10^{23}$	atoms C. 6.02 x molecule		D. 6.02 x 10^{23} entities	
112. What is the weight of one molect	le of a monostomic	element X who	ose stomic weight is 36?	
A. $6.0 \ge 10^{-23}$ gm B. $6.02 \ge 10^{23}$			D. $36 \times 10^{-23} \text{ gm}$	
	8	· 8		
113. When α -particles are set through a thin metal foil, most of them go straight through the foil because				
A. α -particles are much heavier than	-	-		
C. α -particles move with high velocit	y D. α -pai	ticles move wit	h low velocity	
114. The reaction, which proceeds in	he forward direction	n, is		
A. $Fe_2O_3 + 6HCl \rightarrow 2FeCl_3 + 3H_2O$			→ NH₄Cl + NaOH	
C. $SnCl_4 + Hg_2Cl_2 \rightarrow SnCl_2 + 2HgCl_2$	D. 2CuI	$+ I_2 + 4K \rightarrow 2Ci$	$u^+ + 4KI$	

115. The first order constant for the decomposition of N_2O_5 is 6.2 x 10 ⁻⁴ sec ⁻¹ . The half-life period for this decomposition in second is				
A. 1117.7	B. 111.7	C. 223.4	D. 160.9	
	amount of zinc is treated s plumes of H_2 evolved is	separately with excess of	2 H ₂ SO ₄ and excess of	
A. 1 : 1	B. 1 : 2	C. 2 : 1	D. 9 : 4	
117. Calcium does no	ot combine directly with			
A. oxygen	B. nitrogen	C. hydrogen	D. carbon	
	rom other elements of its s)	
A. availability of d-o	-		o-ordination number four	
C. its tendency to cat	enate	D. its unique ability to	form multiple bonds	
119. Iodine reacts wi	th cold dil. NaOH to give	0		
A. NaI + $H_2O + O_2$	Ũ	C. NaI + NaIO + H_2O	D. NaI + NaIO ₃ + H_2O	
			и о '	
	somers for the atomic con	npound of the formula C		
A. 2	B. 3	C. 4	D. 5	
	(1			
	llowing is not true in linea	r programming problem	?	
A. A column in the	7			
simplex table that contains all of the				
variables in the solution	ion			
is called pivot or key	5			
column.	2.			
B. A basic solution	A.			
which is also in the feasible region is call	ed			
a basic feasible				
solution.				
C. A surplus variable	is			
a variable subtracted from the left hand sid	ام			
of a greater than or				
equal to constraint to				
convert it into an				
equality. D. A slack variable is				
variable added to the	o a			

left hand side of a less than or equal to constraint to convert it into an equality.

122. The equation of the circle whose diameter lies on 2x + 3y = 3 and 16x - y = 4 and which passes through (4, 6) is B. $5(x^2 + y^2) - 4x - 8y = 200$ D. $5(x^2 + y^2) - 3x - 8y = 200$ A. $x^2 + y^2 = 40$ C. $x^2 + y^2 - 4x - 8y = 200$ 123. Let n(A) = 4 and n(B) = 5. The number of all possible injections from A to B is B. 9 A. 120 C. 24 D. none 124. If $aN = \{ax : x \in N\}$ and $bN \cap cN = dN$, where $b, c \in N$ are relatively prime, then A. c = bdB. b = cdC. d = bcD. none of the above 125. A square root of 3 + 4i is A. $\sqrt{3} + i$ B. 2 - i C. 2 + iD. none of the above 126. Which of the following is not applicable for a complex number? B. Division C. D. Subtraction Addition A. Inequality 127. | maximum amp (z) - minimum amp (z) | is equal to A. $\sin^{-1}(3/5) - \cos^{-1}(3/5)$ B. $\pi/2 + \cos^{-1}(3/5)$ D. $\cos^{-1}(3/5)$ C. π - 2 cos ⁻¹ (3/5) 128. If e, e' be the eccentricities of two conics S and S' and if $e^2 + e'^2 = 3$, then both S and S' can be B. ellipses C. parabolas A. hyperbolas D. none of the above 129. A stick of length 'I' rests against the floor and a wall of a room. If the stick begins to slide on the floor, then the locus of its middle point is B. a parabola A. an ellipse C. a circle D. a straight line 130. The eccentricity of the ellipse which meets the straight line x/y + y/2 = 1 on the axis of x and the straight line x/3 - y/5 = 1 on the axis of y and whose axes lie along the axes of coordinates is C. √6/7 A. 2√6/7 B. $3\sqrt{2}/7$ D. none of the above 131. A and B are positive acute angles satisfying the equations $3\cos^2 A + 2\cos^2 B = 4$ and $3\sin^2 B = 4$ $A/\sin B = 2 \cos B/\cos A$, then A + 2B is equal to A. $\pi/3$ B. $\pi/2$ C. $\pi/6$ D. $\pi/4$

132. At a point 15 metres away from the base of a 15 metres high house, the angle of elevation of the top is			
A. 90°	B. 60°	C. 30°	D. 45°
133. If $tan(\pi \cos \theta) = c$	ot($\pi \sin \theta$), $0 < \theta < 3\pi/4$,	then $\sin(\theta + \pi/4)$ equals	
A. 1/√2	B. 1/2	C. 1/(2√2)	D. √2
134. In a triangle ABC, $(\sin \angle BAD)/(\sin \angle CA)$		and D divides BC interna	lly in the ratio1 : 3. Then
A. √2/3	B. 1/√3	C. 1/√6	D. 1/3
135. The straight line 5	x + 4y = 0 passes throug	h the point of intersection	n of the lines
A. $x + y - 2 = 0$, $3x + 4$	y - 7 = 0	B. $x - y = 0, x + y = 0$	
C. $x + 2y - 10 = 0, 2x + 0$	-	D. none of the above	
	j · c · ·	<u>^'</u>	
136. The number of co	mmon tangents of the cir	cles $x^2 + y^2 - 2x - 1 = 0$	and $x^2 + y^2 - 2y - 7 = 0$ is
A. 4	B. 1	C. 3	D. 2
	D . 1		D. 2
137. If the product of the	ne roots of the equation of	$\alpha x^{2} + 6x + \alpha^{2} + 1 = 0$ is -	2. then α equals
A2	B1	C.2	D. 1
	21 1		2
138. If the roots of a_1x^2	$a^2 + b_1 x + c_1 = 0$ and $a_2 x^2$	$+b_2x + c_2 = 0$ are same,	then
A. $a_1/a_2 = b_1/b_2 = c_1/c_2$	C	B. $a_1 = b_1 = c_1$, $a_2 = b_2 = b_2$	= C ₂
		,	-
C. $a_1 = a_2$, $b_1 = b_2$, $c_1 =$	c ₂	D. $c_1 = c_2$	
	.0,		
139. The roots of the ed	quation $(3 - x)^4 + (2 - x)^4$	$= (5 - 2x)^4$ are	
139. The roots of the ec A. two real and two im	quation $(3 - x)^4 + (2 - x)^4$	$= (5 - 2x)^4$ are B. all imaginary	
139. The roots of the ed	quation $(3 - x)^4 + (2 - x)^4$	$= (5 - 2x)^4$ are	
139. The roots of the ec A. two real and two im	quation $(3 - x)^4 + (2 - x)^4$	$= (5 - 2x)^4$ are B. all imaginary	
139. The roots of the ec A. two real and two im	quation $(3 - x)^4 + (2 - x)^4$	$= (5 - 2x)^4$ are B. all imaginary	
139. The roots of the eq A. two real and two im C. all real 140. The value \int_{-10}^{10} (-	quation $(3 - x)^4 + (2 - x)^4$	$= (5 - 2x)^4$ are B. all imaginary	D1
139. The roots of the eq A. two real and two im C. all real 140. The value $\sum_{x=1}^{10}$ (- A. 10	quation $(3 - x)^4 + (2 - x)^4$ aginary 1) ⁿ is B. 0	= (5 - 2x) ⁴ are B. all imaginary D. none of the above C. 1	
139. The roots of the eq A. two real and two im C. all real 140. The value $\sum_{x=1}^{10}$ (- A. 10 141. If the 10th term of	quation $(3 - x)^4 + (2 - x)^4$ aginary 1) ⁿ is B. 0 E a G.P. is 9 and 4th term	 = (5 - 2x)⁴ are B. all imaginary D. none of the above C. 1 is 4, then its 7 th term is 	
139. The roots of the eq A. two real and two im C. all real 140. The value $\sum_{x=1}^{10}$ (- A. 10	quation $(3 - x)^4 + (2 - x)^4$ aginary 1) ⁿ is B. 0	= (5 - 2x) ⁴ are B. all imaginary D. none of the above C. 1	
139. The roots of the eq A. two real and two im C. all real 140. The value $\sum_{x=1}^{10}$ (- A. 10 141. If the 10th term of A. 9/4	quation $(3 - x)^4 + (2 - x)^4$ aginary 1) ⁿ is B. 0 E a G.P. is 9 and 4th term B. 4/9	 = (5 - 2x)⁴ are B. all imaginary D. none of the above C. 1 is 4, then its 7 th term is 	
139. The roots of the equation A. two real and two im C. all real 140. The value of $\sum_{x=1}^{10}$ (- A. 10 141. If the 10th term of A. 9/4 142. 1 - 1/2 + 1/3 - 1/4	quation $(3 - x)^4 + (2 - x)^4$ aginary 1) ⁿ is B. 0 F a G.P. is 9 and 4th term B. 4/9 + to ∞ equals	 = (5 - 2x)⁴ are B. all imaginary D. none of the above C. 1 is 4, then its 7 th term is C. 6 	D. 36
139. The roots of the eq A. two real and two im C. all real 140. The value $\sum_{x=1}^{10}$ (- A. 10 141. If the 10th term of A. 9/4	quation $(3 - x)^4 + (2 - x)^4$ aginary 1) ⁿ is B. 0 E a G.P. is 9 and 4th term B. 4/9	 = (5 - 2x)⁴ are B. all imaginary D. none of the above C. 1 is 4, then its 7 th term is 	
139. The roots of the equation A. two real and two im C. all real 140. The value of $\sum_{x=1}^{10}$ (- A. 10 141. If the 10th term of A. 9/4 142. 1 - 1/2 + 1/3 - 1/4	quation $(3 - x)^4 + (2 - x)^4$ aginary 1) ⁿ is B. 0 F a G.P. is 9 and 4th term B. 4/9 + to ∞ equals	 = (5 - 2x)⁴ are B. all imaginary D. none of the above C. 1 is 4, then its 7 th term is C. 6 	D. 36
139. The roots of the eq A. two real and two im C. all real 140. The value $\sum_{x=1}^{10}$ (- A. 10 141. If the 10th term of A. 9/4 142. 1 - 1/2 + 1/3 - 1/4 A. log 2 143. 9/1! + 19/2! + 35/3	quation $(3 - x)^4 + (2 - x)^4$ aginary 1) ⁿ is B. 0 F a G.P. is 9 and 4th term B. 4/9 + to ∞ equals	 = (5 - 2x)⁴ are B. all imaginary D. none of the above C. 1 is 4, then its 7 th term is C. 6 C. e⁻¹ 	D. 36
139. The roots of the eq A. two real and two im C. all real 140. The value $\sum_{x=1}^{10}$ (- A. 10 141. If the 10th term of A. 9/4 142. 1 - 1/2 + 1/3 - 1/4 A. log 2	quation $(3 - x)^4 + (2 - x)^4$ aginary 1) ⁿ is B. 0 F a G.P. is 9 and 4th term B. 4/9 + to ∞ equals B. e	 = (5 - 2x)⁴ are B. all imaginary D. none of the above C. 1 is 4, then its 7 th term is C. 6 C. e⁻¹ 	D. 36


144. How many different arrangements can be made out of the letters in the expansion $A^2B^3C^4$, when written in full? C. 2! + 3! + 4! (2! 3! D. 2! 3! - 4! A. 9!/(2! + 3! + 4!)B. 9!/(2! 3! 4!) 4!) 145. The number of straight lines that can be drawn out of 10 points of which 7 are collinear is A. 23 B. 21 C. 25 D. 24 146. $1/n! + 1/[2! (n - 2)!] + 1/[4! (n - 4)!] + \dots$ is A. $(2^{n-1})/n!$ B. $2^{n}/[(n+1)!]$ C. $2^{n}/n!$ /[(n - 1)!]147. The term independent of x in $(x^2 - 1/x)^9$ is B. 49 A. 1 C. -1 D. none of the above 148. The 9th term of an A.P. is 499 and 499th term is 9. The term which is equal to zero is A. 501th B. 502th C. 500th D. none of the above 149. If A $\begin{bmatrix} 3 & 4 \\ 2 & 4 \end{bmatrix}$, B = $\begin{bmatrix} -2 & -2 \\ 0 & -1 \end{bmatrix}$ then (A + B) B. A $^{-1}$ + B $^{-1}$ A. is a skew symmetric matrix C. does not exist D. none of the above 150. If AB = A and BA = B, then B^2 is equal to A.B B. A C. 1 D. 0 $\begin{vmatrix} a & b & 2a\alpha + 3b \\ b & c & 2b\alpha + 3c \\ 2a\alpha + 3b & 2b\alpha + 3c & 0 \end{vmatrix} = 0, \text{ then }$ 151. If the determinant

A. a, b, c are in H.P. B. α is a root of $4ax^2 + 12bx + 9c = 0$ or a, b, c are in G.P. C. a, b, c are in G.P. only a, b, c are in A.P. 152. The value of K so that (x - 1)/-3 = (y - 2)/2K = (z - 3)/2 and (x - 1)/3K = (y - 1)/1 = (z - 6)/-35 may be perpendicular is given by C. -10 D. 10/7 A. -7/10 B. -10/7

153. The equation of the plane containing the line \rightarrow endra + 2j - k) = 0B. + 2j - k) = 3C. i

D. none of the above

154. The mean of discrete observations y_1 , y_2 , y_n is given by



155. For a poisson distribution whose mean is λ , the standard deviation will be A. λ^2 C. $\sqrt{\lambda}$ B. $1/\lambda$ D. λ

156. If a, b, c, d are constants such that a and c are both negative and r is the correlation coefficient between x and y, then the correlation coefficient between $(ax + b)$ and $(cy + d)$ is equal to					
A. (a/c)r	B. c/a	C r	D. r		
-	card from a pack of 52 p til he draws a spade, the B. 9/16		and shuffles the pack. He n the first two draws is D. 1/64		
158. In tossing 10 coins A. 193/256	s, the probability of getti B. 9/128	ng exactly 5 heads is C. 1/2	D. 63/256		
	ed 00, 01, 10, 11 respecting replaced each time, th 3, is				
A. 100/256	B. 231/256	C. 25/256	D. none of the above		
160.The value $\int_{0}^{\pi/4} tat$	$n^2 x dx$ is equal to				
Α. π/4	B. $1 + (\pi/4)$	C.1 - (π/4)	D. none of the above		
161. Let $f[x + (1/x)] =$	$[x^{2} + (1/x^{2})](x \neq 0)$, then	f(x) is equal to			
A. $x^2 - 1$	$[x^{2} + (1/x^{2})](x \neq 0)$, then B. $x^{2} - 2$	$C. x^2$	D. none of the above		
162. Let $f(x) = [tan(\pi/4 so that it is continous e$	$(x - x)]/\cot 2x, x \neq \pi/4$. The verywhere is	e value which should be	assigned to f at $x = \pi/4$,		
A. 1	B. 1/2	C. 2	D. none of the above		
$f_2(x)$ is	re defined on domains D	D_1 and D_2 respectively, the	en domain of $f_1(x)$ +		
A. $D_1 \cap D_2$	$\mathbf{B}.\mathbf{D}_1 \cup \mathbf{D}_2$	C. D ₁ - D ₂	D. D ₂ - D ₁		
164. The derivative of sin x^3 with respect to $\cos x^3$ is equal to					
A $\tan x^3$	B $\cot x^3$	C. $\cot x^3$	D. $\tan x^3$		
165. If $y = f(x)$ is an odd differentiable function defined on (∞, ∞) such that $f'(3) = -2$, then $f'(-3)$ equals					
A. 4	B. 2	C2	D. 0		
166. The line (x/a) + (y A. (a, ba)	t/b) = 1 touches the curve B. (a, a/b)	$y = be^{-x/a}$ at the point C. (a, b/a)	D. none of the above		

167. The least value of 'a' for which the equation $(4/\sin x) + [1/(1 - \sin x)] = a$ has at least one solution on the interval $(0, \pi/2)$ is					
A. 4	B. 1	C. 9	D. 8		
168. The area bounded	by the curve $y^2 = 8x$ and	$1 x^2 = 8y$ is			
A. 32/7	B. 24/5	C. 72/3	D. 64/3		
169. The integrating factor of the differential equation $[(dy/dx)(x \log x)] + y = 2 \log x$ is given by					
A. $\log(\log x)$	B. e^x	C. log x	D. x		
170. If $y = \tan^{-1}[(\sin x + \cos x)/(\cos x - \sin x)]$, then dy/dx is equal to					
A. 1/2	B. 0	C. 1	D. none of the above		
171. The length of tangent from (5, 1) to the circle $x^2 + y^2 + 6x - 4y - 3 = 0$ is					
A. 81	B. 29	C. 7	D. 21		
172. The equation of the straight line which is perpendicular to $y = x$ and passes through (3, 2) will be given by					
A. $x - y = 5$	B. $x + y = 5$	C. $x + y = 1$	D. x - y = 1		
173. If the imaginary part of $(2z + 1)/(iz + 1)$ is - 2, then the locus of the point representing z in the complex plane is					
A. a circle	B. a straight line	C. a parabola	D. none of the above		
174. The sum of 40 terms of an A.P. whose first term is 2 and common difference 4, will be					
A. 3200	B. 1600	C. 200	D. 2800		
175. If a, b, c are in A.P., then a/bc, 1/c, 2/b are in					
A. A.P.	B. G.P.	С. Н.Р.	D. none of the above		
176. The term independent of x in $[x^2 + (1/x^2)]$ is					
A. 1	B 1	C. 48	D. none of the above		
177. The equation of a line through $(2, -3)$ parallel to y-axis isA. $y = -3$ B. $y = 2$ C. $x = 2$ D. $x = -3$					
178. The value $\int_{-2}^{2} (ax^3 + bx + c) dx$ depends of $\int_{-2}^{2} on$					
A. the value of b	B. the value of c	C. the value of a	D. the value of a and b		

179. The range of the function $f(x) = (1 + x^2)/x^2$ is equal to				
A. [0, 1]	B. [1, 0]	C. (1, ∞)	D. [2, ∞]	

180. Two vectors are said to be equal if

A. their magnitudes are same

C. they meet at the same point

B. direction is same

D. they have magnitude and same sense of direction

gr